

Noether conservation laws issue from the gauge invariance of an Euler–Lagrange operator, but not a Lagrangian

G.Sardanashvily

Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

E-mail: sard@grav.phys.msu.su

URL: <http://webcenter.ru/~sardan/>

Abstract. As is well known, there are different Lagrangians which lead to the same Euler–Lagrange operator. The gauge invariance of a Lagrangian guarantees that of the corresponding Euler–Lagrange operator, but not *vice versa*. We show that the gauge invariance of an Euler–Lagrange operator, but not a Lagrangian results in Noether conservation laws.

Let us consider a first order field theory on a smooth fibre bundle $Y \rightarrow X$ coordinated by (x^λ, y^i) . Its configuration space is the first order jet manifold J^1Y of sections of $Y \rightarrow X$ equipped with the adapted coordinates $(x^\lambda, y^i, y_\mu^i)$ (see, e.g., [2, 3]). We will use the notation $\omega = d^n x$, $n = \dim X$, and $\omega_\lambda = \partial_\lambda \rfloor \omega$.

A first order Lagrangian is a density

$$L = \mathcal{L}(x^\lambda, y^i, y_\lambda^i) \omega : J^1Y \rightarrow \wedge^n T^*X \quad (1)$$

on J^1Y . Its variation δL is the second order Euler–Lagrange operator

$$\begin{aligned} \mathcal{E}_L : J^2Y &\rightarrow T^*Y \wedge (\wedge^n T^*X), \\ \mathcal{E}_L &= (\partial_i \mathcal{L} - d_\lambda \partial_i^\lambda) \mathcal{L} \theta^i \wedge \omega, \end{aligned} \quad (2)$$

where $\theta^i = dy^i - y_\lambda^i dx^\lambda$ are contact forms and $d_\lambda = \partial_\lambda + y_\lambda^i \partial_i + y_{\lambda\mu}^i \partial_i^\mu$ are the total derivative. Its kernel $\text{Ker } \mathcal{E}_L \subset J^2Y$ defines the Euler–Lagrange equations. There are different Lagrangians whose variations provide the same Euler–Lagrange operator. They make up an affine space modelled over the vector space of variationally trivial Lagrangians L_0 , i.e., $\delta L_0 = 0$. One can show that a first order Lagrangian L_0 is variationally trivial iff

$$L_0 = h_0(\varphi), \quad (3)$$

where φ is a closed n -form on Y and

$$h_0(dx^\lambda) = dx^\lambda, \quad h_0(\theta^i) = 0$$

is the horizontal operator acting on semibasic exterior forms on $J^1Y \rightarrow Y$.

Gauge transformations are defined as bundle automorphisms of a fibre bundle $Y \rightarrow X$. Any projectable vector field

$$u = u^\lambda(x^\mu)\partial_\lambda + u^i(x^\mu, y^j)\partial_i \quad (4)$$

on $Y \rightarrow X$ is the infinitesimal generator of a local one-parameter group G_u of gauge transformations of $Y \rightarrow X$, and *vice versa*. Its jet prolongation onto J^1Y reads

$$J^1u = u^\lambda\partial_\lambda + u^i\partial_i + (d_\lambda u^i - y_\mu^i\partial_\lambda u^\mu)\partial_i^\lambda. \quad (5)$$

A Lagrangian L (1) is invariant under a gauge group G_u iff its Lie derivative

$$\mathbf{L}_{J^1u}L = [\partial_\lambda(u^\lambda\mathcal{L}) + (u^i\partial_i + (d_\lambda u^i - y_\mu^i\partial_\lambda u^\mu)\partial_i^\lambda)\mathcal{L}]\omega \quad (6)$$

along J^1u vanishes.

The first variational formula of the calculus of variations provides the canonical decomposition

$$\mathbf{L}_{J^1u}L = u_V]\mathcal{E}_L + h_0d(u]H_L) \quad (7)$$

of the Lie derivative (6), where: $u_V = (u]\theta^i)\partial_i$, $\mathcal{E}_L$ is the Euler–Lagrange operator (2) and H_L is some Lepagean equivalent of L , e.g., the Poincaré–Cartan form

$$H_L = L + \partial_i^\lambda \mathcal{L} \theta^i \wedge \omega_\lambda = \partial_i^\lambda \mathcal{L} dy^i \wedge \omega_\lambda + (\mathcal{L} - y_\lambda^i \partial_i^\lambda \mathcal{L})\omega.$$

For instance, if $L = L_0$ is a variationally trivial Lagrangian, the first variational formula (7) gives the equality

$$\mathbf{L}_{J^1u}L_0 = h_0d(u]H_{L_0}). \quad (8)$$

If the Lie derivative (6) vanishes, one obtains the weak conservation law

$$0 \approx h_0d(u]H_L) = d_\lambda[\pi_i^\lambda(u^\mu y_\mu^i - u^i) - u^\lambda\mathcal{L}]\omega \quad (9)$$

on $\text{Ker } \mathcal{E}_L$.

The Euler-Lagrange operator $\mathcal{E}_L$ (2) is invariant under a one-parameter gauge group G_u iff its Lie derivative $\mathbf{L}_{J^2u}\mathcal{E}_L$ along the jet prolongation J^2u of u onto the second order jet manifold J^2Y vanishes. There is the relation

$$\mathbf{L}_{J^2u}\mathcal{E}_L = \delta(\mathbf{L}_{J^1u}L) = \mathcal{E}_{\mathbf{L}_{J^1u}L}, \quad (10)$$

i.e., the Lie derivative $\mathbf{L}_{J^2u}\mathcal{E}_L$ is the Euler–Lagrange operator associated to the Lagrangian $\mathbf{L}_{J^1u}L$ [1, 2]. It follows that $\mathbf{L}_{J^2u}\mathcal{E}_L = 0$ iff the Lie derivative $\mathbf{L}_{J^1u}L$ is a variationally trivial Lagrangian, i.e., is given by the expression (3). Substituting this expression into the first variational formula (7), we obtain the weak equality

$$h_0(\varphi) \approx h_0d(u \rfloor H_L) \quad (11)$$

on $\text{Ker } \mathcal{E}_L$. If $\varphi = d\sigma$ is an exact form on Y , this equality is brought into the weak conservation law

$$0 \approx h_0d(u \rfloor H_L - \sigma). \quad (12)$$

Let $L' = L + L_0$ be another Lagrangian whose variation is the Euler–Lagrange operator $\mathcal{E}_L$. Due to the equality (8), we come to the same conservation law (12).

References

- [1] G.Giachetta and L.Mangiarotti, Gauge invariant and covariant operators in gauge theories, *Int. J. Theor. Phys.* **29** (1990) 789.
- [2] G.Giachetta, L.Mangiarotti and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field Theory* (World Scientific, Singapore, 1997).
- [3] G.Sardanashvily, Ten lectures on jet manifolds in classical and quantum field theory, *E-print arXiv: math-ph/0203040*.